



CITY OF LOS ANGELES

2024 COMMUNITY GREENHOUSE GAS INVENTORY REPORT



LACITYSAN.ORG/CLIMATEACTION



SAN.CLIMATEACTION@LACITY.ORG



Table of Contents

Executive Summary	1
1. Introduction	3
2. Methodology	4
2.1 Scopes.....	4
2.2 BASIC vs BASIC+	5
2.3 Data Collection and Providers.....	6
3. Findings by Sector	7
3.1 Stationary Energy	8
3.2 Transportation.....	11
3.3 Waste	14
3.4 Agriculture, Forestry, and Other Land Use	17
3.5 Industrial Processes and Product Use	19
4. Conclusion	21
5. Preparers	22

List of Figures

Figure 1. Emissions Progress Compared to Targets.....	1
Figure 2. Emissions Trends and Metrics.....	2
Figure 3. Sources Included in BASIC and BASIC+	5
Figure 4. Basic Emissions by Sector	7
Figure 5. BASIC Stationary Energy Sector Emissions Reductions from Baseline	9
Figure 6. Electricity Generation Portfolio vs Emissions	9
Figure 7. BASIC Transportation Sector Emissions Reductions from Baseline	12
Figure 8. BASIC Waste Sector Emissions Reductions from Baseline	15
Figure 9. Landfilled Waste Characterization vs. Emissions.....	16
Figure 10. AFOLU Sector Emissions Reductions from Baseline.....	18
Figure 11. IPPU Sector Emissions Compared to Baseline.....	20

List of Tables

Table 1. BASIC Emissions by Sector (Million Metric Tons CO ₂ e)	2
Table 2. Greenhouse Gas Global Warming Potential Factors.....	4
Table 3. Community GHG Inventory Data Providers.....	6
Table 4. BASIC Stationary Energy Emissions by Subsector (MT CO ₂ e)	8
Table 5. BASIC+ Stationary Energy Emissions by Subsector (MT CO ₂ e).....	10
Table 6. BASIC Transportation Emissions by Subsector (MT CO ₂ e)	11
Table 7. BASIC+ Transportation Emissions by Subsector (MT CO ₂ e)	12
Table 8. Waste Emissions by Subsector (MT CO ₂ e).....	14
Table 9. AFOLU Emissions by Subsector (MT CO ₂ e).....	17
Table 10. IPPU Emissions by Subsector (MT CO ₂ e)	19

Executive Summary

In 2024, Los Angeles achieved a significant milestone in its environmental efforts, with community-wide greenhouse gas (GHG) emissions at 31% below the 1990 baseline, as depicted in Figure 1. The City is making progress towards its interim targets of a 50% reduction by 2025 and a 73% reduction by 2035. These efforts are part of a broader, ambitious plan encapsulated in Los Angeles' Green New Deal, which sets the ultimate goal of reaching carbon neutrality by 2050. This commitment reflects the City's proactive approach to mitigating climate change and fostering a sustainable future.




Figure 1. Emissions Progress Compared to Targets

Throughout the 2024 Inventory, several data sources that were usually used were unavailable or behind schedule in their updates, resulting in the use of scaled data from other sources, trends from previous years, or placeholders in lieu of the usual sources. Report text will indicate areas where this is significant and updates to this report or the values in future years' Inventory Reports will include revised values once available.

In 2024, Los Angeles experienced a slight increase in community-wide GHG emissions compared to 2023, primarily from the transportation sector. Total emissions for the year amounted to 26.1 million metric tons of carbon dioxide equivalent (MMT CO₂e), for an increase of about 0.8 million metric tons compared to the total in 2023. This leaves a gap of 7.3 MMT CO₂e to achieve the target of reducing emissions to 50% below the 1990 baseline in 2025.

Table 1. BASIC Emissions by Sector (Million Metric Tons CO₂e)

	1990	2018	2019	2020	2021	2022	2023	2024	1990 vs 2024 Percent Change
Stationary Energy	26.0	19.0	17.7	16.8	16.8	15.6	14.6	14.5	-44%
Transportation	10.4	11.1	10.7	7.4	9.0	9.3	9.2	10.1	-3%
Waste	1.2	1.3	1.3	1.3	1.2	1.3	1.5	1.5	25%
Total Emissions	37.6	31.3	29.8	25.6	27.0	26.2	25.3	26.1	-31%

As detailed in Table 1, the stationary energy sector has seen a substantial decrease of 44% in emissions compared to the 1990 baseline, declining to 14.5 MMT CO₂e in 2024. The transportation sector had 10.1 MMT CO₂e in 2024, placing it 3% below the 1990 levels. However, it is using extrapolated data that is discussed in report section 3.2 Transportation. The waste sector showed an increase of 25% compared to 1990, settling at 1.5 MMT CO₂e which is the same as in 2023 after rounding. Overall, these sectors contributed to a total decrease of 31% in GHG emissions compared to the 1990 baseline.

Figure 2 illustrates that despite an increase in gross domestic product (GDP) from 2023¹, the ratio of emissions per GDP unit has continued its downward trend. This ongoing reduction in emissions relative to economic output effectively demonstrates the decoupling of GHG emissions from economic growth, indicating progress towards a more sustainable and less carbon-intensive economy.




Figure 2. Emissions Trends and Metrics

¹ 2024 data applied a trend from 2001-2023 for the ratio of LA county GDP to CA state GDP: U.S. Bureau of Economic Analysis, "[CAGDP9 Real GDP by county and metropolitan area](#)" (accessed Tuesday, January 13, 2026).

1. Introduction

Climate change, a critical global challenge, requires decisive local actions. The City of Los Angeles (City) is at the forefront of this battle, implementing robust measures to curb greenhouse gas (GHG) emissions and mitigate climate impacts. LA Sanitation and Environment (LASAN) plays a vital role in the City's climate strategy. Understanding the quantity, sources, and trends of Los Angeles' GHG emissions is crucial for effective climate action. This knowledge enables policymakers and stakeholders to take targeted and adaptive measures against climate change. LASAN's annual, detailed community GHG inventories are instrumental in gaining this insight and tracking the City's progress toward its ambitious climate objectives.

Since 2019, as originally requested by the Mayor's Office of Sustainability (MOS), LASAN has been responsible for preparing the City of Los Angeles' annual Community GHG Inventory. This task included enhancing the previously prepared inventories from a BASIC to a BASIC+ rating, a change necessitated by the availability of more comprehensive data sources. The BASIC+ inventory offers a deeper understanding of Los Angeles' emissions scenario because it includes industrial processes, product use, agriculture, forestry, and other land uses, along with some Scope 3 emissions from stationary sources and transportation sectors.

To date, LASAN has compiled community-wide inventories for the years 2014-2024, in addition to the baseline year of 1990, as established in the City's Sustainable City pLAn. This report presents the 2024 Community GHG Inventory values alongside the updated values for the City's prior inventories, offering insights into the City's emission trends over time and using the 1990 figures as the benchmark for all reduction percentages.

2. Methodology

LASAN compiles the City of Los Angeles' Community Greenhouse Gas (GHG) Inventory adhering to the standards set by C40's Global Protocol for Community-Scale Greenhouse Gas Emissions (GPC), a globally recognized framework. This approach integrates two key methods: the scopes framework and the city-induced framework, accommodating emissions generated both inside and outside the City's borders. The inventory accounts for five GHGs: carbon dioxide (CO₂), methane (CH₄), nitrous oxide (N₂O), hydrofluorocarbons (HFCs), and perfluorocarbons (PFCs), each with a specific Global Warming Potential (GWP). GWP measures a gas's heat-trapping ability relative to CO₂. For a consistent assessment, emissions of these gases are converted to CO₂ equivalents (CO₂e) throughout this report.

Table 2. Greenhouse Gas Global Warming Potential Factors

Greenhouse Gas	Formula	GWP*
Carbon Dioxide	CO ₂	1
Methane	CH ₄	25
Nitrous Oxide	N ₂ O	298
Hydrofluorocarbons	HFCs	124-14,800
Perfluorocarbons	PFCs	7,390-12,200
*100-year time horizon.		

This inventory employs the Intergovernmental Panel on Climate Change's (IPCC) AR4 GWP values, aligning with the California Air Resources Board's (CARB) methodology for the statewide California Greenhouse Gas Emissions Inventory (see Table 2).

2.1 Scopes

Scope 1 emissions are from sources located within the City boundary (in-boundary activities). These can also be considered "territorial" emissions because they are all produced within the geographic boundary. Scope 2 emissions occur from the use of grid-supplied electricity, heat, steam, and/or cooling within the city boundary. Scope 3 emissions are from sources outside the city boundary because of actions occurring within the city boundary (out-of-boundary activities). Figure 3 provides examples of all three scopes as used in this inventory. For example, waste entirely produced within the City and disposed of at facilities outside of the City produces emissions when it breaks down. These emissions would still be attributed to the City as Scope 3 emissions, since it came from activity within the City boundary.

2.2 BASIC vs BASIC+

The GPC categorizes two levels, BASIC and BASIC+ (see Figure 3). BASIC encompasses Scopes 1 and 2 emissions from stationary energy and transportation and Scopes 1 and 3 from waste. BASIC+ extends this to include industrial processes and product use (IPPU), as well as agriculture, forestry, and other land use (AFOLU), plus Scope 3 emissions from stationary energy and transportation sectors. Although BASIC+ offers a broader emission perspective, the City has limited direct control over these additional sectors. This report discusses all sectors under BASIC+ but tracks progress towards L.A.'s Green New Deal goals using BASIC emissions.




Figure 3. Sources Included in BASIC and BASIC+

2.3 Data Collection and Providers

LASAN's inventory preparation involves collaboration with multiple departments and agencies, ensuring a comprehensive data collection process. Table 3 summarizes these data contributors.

Table 3. Community GHG Inventory Data Providers

Data Provider	Data
City Department	
LASAN	Solid waste characterization; compost tonnage; biosolids; wastewater treatment
LADWP	Residential, commercial, institutional, and industrial electricity consumption; transmission and distribution losses; water services-related electricity consumption, power generation fuel consumption, EV charging electricity consumption
LAWA	Commercial jet fuel usage
POLA	Local harbor craft fuel usage
LA Animal Services Department	Livestock estimates
Regulatory Agency	
CARB	EMFAC2021 Model for vehicle fuel efficiency; off-road transportation emissions estimate; fuel estimate for vessel bunkering; industrial facilities involved with mineral, chemical, or metal production; Substitutes for Ozone Depleting Substance usage
South Coast Air Quality Management District (SCAQMD)	Industrial fuel consumption, landfill flaring
Environmental Protection Agency	Refinery feed flaring; industrial facilities involved with mineral, chemical, or metal production; eGRID
California Geologic Energy Management (CalGEM)	Oil and gas wells
Federal Air Administration	Local aircraft fuel usage
CalRecycle	Solid waste disposal tonnage
California Department of Food and Agriculture	Fertilizer usage
Other	
Energy Information Administration (EIA)	Residential, commercial, and industrial wood and ethanol consumption estimate
SoCal Gas	Residential, commercial, and industrial natural gas consumption
Southern California Edison	Utility's electricity emission factor
Google Environmental Insights Explorer	On-road VMTs
Amtrak	Annual trips estimate
Metrolink	Annual trips estimate
Burlington Northern Santa Fe (BNSF) Railway	Fuel usage
Union Pacific	Fuel usage
LA Metro	Rail propulsion electricity usage
International Council for Local Environmental Initiatives (ICLEI)	Land use change emissions estimate
US Census	Population
US Bureau of Economic Analysis	County and State GDP

3. Findings by Sector

The following sections present findings from the City's 2014-2024 Community GHG Inventories. Each year, LASAN continues to update the community GHG inventory to incorporate new procedures, as well as make improvements to data collection processes, methodologies, emissions factors, and quality assurance. Inventories from previous years are updated and revised to reflect these changes and to maintain a consistent time series following recommendations from the IPCC for developing GHG inventories. As a result, reported emissions may differ from values published in prior inventory reports.

As shown in Figure 4, the largest sector in Los Angeles' Community GHG Inventory (at the BASIC level) is the stationary energy sector, accounting for 55% of total emissions, followed by the transportation sector and the waste sector. The magnitude of stationary energy emissions continues to decrease, resulting in the transportation and waste sectors representing an increasingly larger proportion of the City's overall community emissions.




Figure 4. Basic Emissions by Sector

Table 1 data displays that between 2014 and 2024, Los Angeles saw significant changes in greenhouse gas (GHG) emissions in key sectors. In the stationary energy sector, emissions decreased from 23.8 million metric tons in 2014 to 14.5 million in 2024, a substantial reduction reflecting the City's energy efficiency and sustainable energy initiatives to reduce usage of fossil fuels. The transportation sector, identified as the second-largest contributor to emissions at the BASIC level, has shown a gradual decline compared to the sector's 1990 baseline but is only 3% below that baseline in 2024². Waste sector emissions, while a smaller portion of the total, were very similar to 2023 values, accounting for about 6% of the City's emissions throughout this period. These trends demonstrate the efficacy of Los Angeles' strategies in reducing its environmental impact in major emission-generating sectors.

² 2024 Transportation on road activity data is not up to date in Google Environmental Insights Explorer. See 3.2 Transportation

3.1 Stationary Energy

The stationary energy sector includes fuel combustion and fugitive emissions that occur while generating, delivering, and consuming useful forms of energy (such as electricity or heat). The five main subsectors are residential buildings, commercial and institutional buildings and facilities, manufacturing industries and construction, energy industries³, and fugitive emissions from oil and natural gas systems.

Figure 5 shows overall emissions in this sector have decreased by 44% since 1990 with reductions primarily driven by decarbonization of the electricity grid in the residential buildings and commercial and institutional buildings and facilities subsectors.

Between 2014 and 2024, the ratio of MWh generated to the GHG emissions of the City's electricity has decreased by 52% (Figure 6).

Table 4. BASIC Stationary Energy Emissions by Subsector (MT CO₂e)

	Residential buildings	Commercial and institutional buildings and facilities	Manufacturing industries and construction	Energy industries	Fugitive emissions from oil and natural gas systems	Total Stationary Energy Emissions
1990	7,188,792	9,903,709	4,297,840	4,146,575	437,185	25,974,102
2014	6,585,035	8,460,163	3,995,230	4,587,057	214,610	23,842,094
2015	6,543,251	8,218,274	3,918,539	4,161,174	213,396	23,054,634
2016	5,394,410	6,226,515	3,446,961	4,294,149	216,983	19,579,017
2017	5,212,031	5,885,760	3,817,250	4,437,001	215,680	19,567,722
2018	5,099,756	5,902,502	3,324,850	4,494,205	216,611	19,037,923
2019	5,023,171	5,536,819	2,724,736	4,219,939	216,173	17,720,837
2020	5,039,695	4,700,410	2,660,925	4,201,503	211,973	16,814,505
2021	4,848,815	4,896,054	2,706,020	4,131,840	208,662	16,791,392
2022	4,641,986	4,856,690	1,840,049	4,061,659	207,544	15,607,928
2023	4,526,202	4,650,203	997,187	4,247,210	202,996	14,623,798
2024	4,330,196	4,621,478	935,424	4,372,985	197,694	14,457,778

³ Energy Industries emissions includes heat plants, petroleum refining, and fuel consumption for oil and gas production auxiliary power. According to GPC, energy industries does not include electricity supplied to the grid in BASIC/BASIC+ reporting. Electricity supplied to the grid is tabulated and tracked separately for internal record keeping of territorial emissions.




Figure 5. BASIC Stationary Energy Sector Emissions Reductions from Baseline




Figure 6. Electricity Generation Portfolio vs Emissions⁴

⁴ Power generation and emissions data provided by LADWP.

LADWP progresses towards supplying 100% carbon-free energy by 2035, as outlined in the LA100 Plan. The City remains steadfast in advancing its ambitious clean energy targets. While significant progress has been made, reaching these goals will require further action to decarbonize buildings, improve energy efficiency across all sectors, and reduce fuel combustion in industrial activities. Notably, the increased emissions observed between the BASIC level in Table 4 and the BASIC+ level in Table 5 are largely attributed to electricity transmission and distribution losses. Including both BASIC and BASIC+ metrics allows us to track emissions within the City's direct influence while also gaining a broader understanding of indirect impacts, such as transmission losses and industrial activities.

In 2024, the stationary energy sector experienced notable trends. Emissions from residential and commercial buildings continued their steady decline. However, emissions from energy industries saw a slight increase, highlighting ongoing challenges in decarbonizing this subsector, while fugitive emissions from oil and natural gas systems showed only a modest decrease.

In the 2023 Community GHG Inventory Report, the manufacturing and construction industries subsector had a substantial drop that was under further investigation. In the data for 2024 this reduction was found to be consistent.

Table 5. BASIC+ Stationary Energy Emissions by Subsector (MT CO₂e)

	Residential buildings	Commercial and institutional buildings and facilities	Manufacturing industries and construction	Energy industries	Fugitive emissions from oil and natural gas systems	Total Stationary Energy Emissions
1990	7,636,543	11,105,676	4,508,456	4,146,575	437,185	27,834,436
2014	7,199,130	9,659,486	4,132,757	4,587,057	214,610	25,793,040
2015	7,100,858	9,311,586	4,031,961	4,161,174	213,396	24,818,912
2016	5,774,009	6,971,781	3,522,091	4,294,149	216,983	20,779,013
2017	5,559,937	6,440,128	3,886,707	4,437,001	215,680	20,539,454
2018	5,428,269	6,564,167	3,379,549	4,494,205	216,611	20,082,800
2019	5,316,558	6,079,030	2,770,620	4,219,939	216,173	18,602,319
2020	5,334,948	4,851,102	2,701,097	4,201,503	211,973	17,300,622
2021	5,130,246	5,454,444	2,744,493	4,131,840	208,662	17,669,685
2022	4,913,838	5,483,767	1,879,337	4,061,659	207,544	16,546,145
2023	4,762,585	5,077,534	1,030,098	4,247,210	202,996	15,320,422
2024	4,561,341	4,994,172	966,916	4,372,985	197,694	15,093,109

3.2 Transportation

The transportation sector includes GHG emissions from fuel combustion and electricity used for transportation activities. The sector covers five subsectors: on-road transportation, railways, waterborne navigation, aviation, and off-road transportation. In 2024, the presented on-road data is linearly extrapolated from data from 2021-2023 because Google Environmental Insights Explorer has not updated values for 2024. Because of the very small sample size and impacts of COVID-19 on the sector in 2020 and 2021, there is the potential for the true values to be significantly different than what is reported here.

Table 6. BASIC Transportation Emissions by Subsector (MT CO₂e)

	On-road transportation	Railways	Waterborne navigation	Aviation	Off-road transportation	Total Transportation Emissions
1990	10,197,626	24,508	36,204	11,833	160,255	10,430,427
2014	11,128,431	98,421	55,892	17,336	239,507	11,539,586
2015	10,980,018	99,356	61,013	16,792	300,569	11,457,748
2016	10,827,904	82,100	58,348	14,906	276,693	11,259,952
2017	10,750,193	83,800	62,331	17,959	224,603	11,138,885
2018	10,605,673	85,269	66,092	16,828	321,258	11,095,119
2019	10,296,705	86,196	60,884	16,157	328,534	10,788,476
2020	6,961,453	76,900	60,734	14,068	327,406	7,440,562
2021	8,486,904	68,835	53,521	15,565	336,063	8,960,887
2022	8,836,599	66,992	50,811	15,116	342,143	9,311,662
2023	8,688,602	67,635	51,808	15,007	349,552	9,172,605
2024	9,638,289	70,679	55,604	14,737	365,831	10,145,141

Overall, transportation sector emissions have decreased by 3% since 1990 (Figure 7), primarily driven by reductions in on-road transportation emissions. The data in Table 6, spanning 2014 to 2024, highlights key trends. Railways have seen a slight increase in emissions between 2022 and 2024, and so have waterborne navigation emissions from an increasing amount of container throughput.

Off-road transportation emissions, which originate from equipment and vehicles operating off public roads—such as construction machinery, agricultural equipment, and recreational vehicles—continue to rise in Los Angeles. Addressing these emissions requires targeted strategies, including transitioning to electric off-road equipment, decarbonizing the grid, and implementing policies that promote cleaner technologies in these sectors. Complementary efforts to reduce on-road vehicle miles traveled (VMTs) through investments in public transportation and walkable, people-centric infrastructure remain critical to achieving broader emission reduction goals.




Figure 7. BASIC Transportation Sector Emissions Reductions from Baseline

The BASIC+ transportation emissions, shown in Table 7, reflect the inclusion of harbor and airport tenant commercial cargo ship and aircraft fuel usage, leading to much higher totals. This broader scope ensures a comprehensive understanding of transportation emissions, balancing the focus on areas the City can influence directly while addressing the global impact of indirect sources. Despite this, the City of Los Angeles is making significant progress in reducing overall transportation emissions, with an 11% decrease since 1990. Efforts to address emissions from aviation and waterborne navigation, such as Los Angeles World Airports' (LAWA) promotion of sustainable aviation fuels and the Port of Los Angeles' (POLA) shore-side electricity program, highlight the City's commitment to tackling emissions even in sectors outside its direct control. These initiatives, combined with ongoing investments in clean transportation and infrastructure, demonstrate the City's dedication to achieving its climate goals.

Table 7. BASIC+ Transportation Emissions by Subsector (MT CO_{2e})

	On-road transportation	Railways	Waterborne navigation	Aviation	Off-road transportation	Total Transportation Emissions
1990	10,197,626	74,287	8,786,351	14,431,582	160,255	33,650,100
2014	11,130,442	205,088	2,866,564	15,837,678	162,034	30,201,806
2015	10,983,242	201,855	3,494,079	17,009,219	220,511	31,908,906
2016	10,834,837	181,741	4,479,269	18,323,514	276,693	34,096,055
2017	10,757,820	188,836	3,043,868	19,147,217	224,603	33,362,344
2018	10,615,057	197,677	2,311,149	19,559,534	321,258	33,004,676
2019	10,306,160	212,229	3,044,750	19,401,839	328,534	33,293,511
2020	6,969,168	187,003	1,898,903	11,451,145	327,406	20,833,625
2021	8,496,451	175,178	2,230,736	13,987,150	336,063	25,225,578
2022	8,847,752	175,258	2,082,060	15,219,042	342,143	26,666,254
2023	8,701,565	163,387	2,546,545	13,328,490	349,552	25,089,538
2024	9,654,595	172,839	2,664,110	17,112,820	365,831	29,970,196

3.3 Waste

The waste sector is responsible for emissions stemming from the disposal and treatment of both solid waste and wastewater, with processes such as decomposition and incineration as primary contributors.

Table 8. Waste Emissions by Subsector (MT CO₂e)⁵

	Solid waste generated in the city	Biological waste generated in the city	Incinerated and burned waste generated in the city	Wastewater generated in the city	Total Waste Emissions
1990	1,120,499	3,642	11,183	47,775	1,183,101
2014	1,046,584	5,798	14,670	44,841	1,111,893
2015	1,111,432	6,871	13,314	46,839	1,178,456
2016	1,157,181	11,692	15,581	44,471	1,228,924
2017	1,228,286	5,189	3,890	49,616	1,286,980
2018	1,262,352	9,454	2,596	51,317	1,325,719
2019	1,267,111	8,363	2,262	53,344	1,331,079
2020	1,256,478	6,650	3,451	49,154	1,315,733
2021	1,186,929	8,213	3,158	48,453	1,246,753
2022	1,250,175	6,672	19,202	49,859	1,325,907
2023	1,413,797	5,626	11,896	49,873	1,481,192
2024	1,421,704	5,626	1,569	51,529	1,480,429

Landfill disposal, primarily from solid waste, remains the largest contributor to emissions within the waste sector, accounting for over 95% of the total (see Table 8). In 2024, solid waste generation in the city increased by 25% compared to 1990, reflecting a rise in consumer activity and shopping trends among Angelenos. However, waste emissions require additional context to avoid misinterpretation. While waste emissions in 2024 may be 25% higher than the 1990 baseline, they represent only 6% of the City's total BASIC emissions. This underscores the relatively minor contribution of waste compared to other sectors, such as stationary energy and transportation, which together account for 94% of emissions.

Additionally, growth in population and consumption since 1990 provides important context for changes in waste-sector emissions. Continued investment in public education, behavioral change programs, and community engagement can support further reductions over time.

⁵ For the waste sector, BASIC and BASIC+ emissions are the same (see Figure 3).




Figure 8. BASIC Waste Sector Emissions Reductions from Baseline

Figure 9, from a 2016 waste characterization study, shows how waste composition significantly affects emissions levels; the different materials have varying organic content that decomposes at different rates. Paper and cardboard make up 23% of the City's waste by volume, yet they contribute to 53% of emissions from landfilled waste. Organic waste, while constituting 26% of the waste stream, accounts for 37% of emissions. These two categories alone represent 90% of emissions from landfilled waste, underscoring the importance of targeted waste reduction in these areas.⁶

⁶ Waste characterization obtained from *Sunshine Canyon Landfill Comprehensive Waste Characterization Study*, May 2016.




Figure 9. Landfilled Waste Characterization vs. Emissions

*Percentages may not total to 100 percent due to rounding.

**Other includes rubber and leather, electronics, gypsum board, inert material, household hazardous waste, special waste, and mixed residue.

3.4 Agriculture, Forestry, and Other Land Use

The AFOLU sector is characterized by GHG emissions from several sources, including land-use changes, enteric fermentation in livestock, and agricultural nutrient management quantities that are aggregated together. According to GPC, in the City's emissions framework, these emissions are detailed only in the BASIC+ inventory, not the BASIC total. While carbon sequestration data is captured for reference, it isn't factored into the net emissions for BASIC+.

Table 9. AFOLU Emissions by Subsector (MT CO₂e)

	Livestock	Land		Aggregate Emission Sources		Total (sources)
	Sources	Sources	Sinks	Sources	Sinks	
1990	4,418	13,981	(112,441)	9,144	(48,556)	27,543
2014	4,418	12,165	(112,706)	5,616	(79,865)	22,198
2015	4,418	12,165	(112,736)	5,689	(80,407)	22,271
2016	4,418	12,165	(112,541)	5,506	(63,510)	22,089
2017	4,418	11,581	(112,797)	6,018	(34,888)	22,016
2018	4,418	11,492	(112,827)	7,517	(45,678)	23,427
2019	4,418	11,344	(112,523)	32,810	(72,411)	48,572
2020	4,418	11,286	(112,729)	8,681	(76,818)	24,384
2021	4,275	11,224	(112,609)	12,144	(66,524)	27,643
2022	4,275	11,115	(112,716)	28,211	(58,158)	43,601
2023	4,275	16,546	(108,509)	3,684	(62,130)	24,505
2024	4,275	18,384	(107,487)	4,556	(65,959)	27,215

The primary reported emissions source in the aggregate emissions sources subsector is synthetic fertilizer use, with a notable increase in 2019 and 2022 emissions driven by greater application of synthetic fertilizers (Figure 10). This Value is scaled down from reported County level data and may not reflect conditions within the City boundary. Initiatives like organic waste diversion and anaerobic digesters have likely contributed to increased compost and biosolids production and application, potentially reducing reliance on synthetic fertilizers and promoting soil carbon sequestration. Beyond reducing emissions, composting offers co-benefits such as water conservation, improved soil microbial health, erosion control, and air quality improvements. This progress highlights the critical role of sustainable waste management practices in mitigating climate impacts and fostering healthier ecosystems.

In 2024, while emissions from land sources increased—potentially due to shifts in land use or reduced sequestration capacity—the City's efforts to scale composting and soil health improvements underscore the potential for targeted programs to drive reductions in greenhouse gas emissions.




Figure 10. AFOLU Sector Emissions Reductions from Baseline

3.5 Industrial Processes and Product Use

The IPPU sector captures emissions from non-energy related industrial processes⁷ and the use of products not related to energy generation. This includes emissions from substitutes for ozone-depleting substances (ODS), such as hydrofluorocarbons (HFCs) and perfluorocarbons (PFCs). Both of these ODS substitutes replaced ozone-depleting Chlorofluorocarbons (CFCs) that were in use in 1990 and these substitutes are now widely used in refrigeration, air conditioning, foam production, and fire suppression systems instead of CFCs. These emissions are part of the BASIC+ standard and thus are not included in the BASIC emissions tally.

Table 10. IPPU Emissions by Subsector (MT CO₂e)

	Industrial Processes	Product Use	Total IPPU Emissions
1990	-	9,753	9,753
2014	-	1,790,652	1,790,652
2015	-	1,879,750	1,879,750
2016	-	1,947,186	1,947,186
2017	-	2,009,201	2,009,201
2018	-	2,050,501	2,050,501
2019	-	2,067,530	2,067,530
2020	-	2,071,863	2,071,863
2021	-	2,061,848	2,061,848
2022	-	2,058,893	2,058,893
2023	-	2,086,144	2,086,144
2024	-	2,106,905	2,106,905

A substantial portion of the IPPU emissions stems from the use of HFCs and PFCs. To mitigate this, the City is exploring policy options to incentivize the adoption of alternative cooling technologies, like cool roofs and cool pavements, which offer the dual benefits of reducing heat absorption and lowering cooling demands. The California Air Resources Board (CARB) is also proactive in this area, developing regulations that encourage the use of substances with a lower climate change potential.

Given the vast technological and industrial changes since, the baseline year for IPPU emissions provides limited insights for comparison. The sector has evolved, with new processes and technologies emerging that were not present three decades ago, which now contribute to GHG emissions. For instance, emissions from IPPU in 1990 were recorded at a mere 10,000 MT CO₂e, contrasting sharply with the over 2 million MT CO₂e reported in recent years.

Thus, while historical data offer context, the focus must be on current and future actions. Efforts to find alternatives to substitutes for ozone-depleting substances are paramount. This includes

⁷ Industrial process facilities are in the mineral, chemical, and metal production industries. No industrial facilities within Los Angeles meet the thresholds for reporting to statewide and national regulatory agencies, including CARB and the US EPA. As a result, industrial process emissions are not in this inventory. This does not necessarily mean there are no industrial process emissions, only that there are no facilities that meet the reporting thresholds.

identifying new chemicals or methods that serve similar functions or devising innovative strategies to reduce the need for refrigerants and cooling agents altogether.

The IPPU sector's evolution underscores the necessity for a dynamic and responsive approach to environmental management, one that acknowledges past practices while aggressively pursuing current opportunities for emission reductions.




Figure 11. IPPU Sector Emissions Compared to Baseline

4. Conclusion

Los Angeles' unwavering determination and strategic action in addressing climate change are exemplified by the substantial greenhouse gas (GHG) emissions reductions recorded in the 2024 Community GHG Inventory. With emissions now 31% below the 1990 baseline, the City continues to demonstrate its commitment to a sustainable future while setting a global benchmark for urban climate leadership.

To meet the ambitious target of a 50% reduction by 2025, Los Angeles must achieve an additional 19% reduction in emissions in the next year's data. This pressing challenge underscores the need for accelerated efforts across all sectors, including scaling clean energy adoption, electrifying transportation systems, enhancing waste reduction and recycling programs, and addressing fugitive emissions. Achieving this goal will require innovation, collaboration, and an unwavering focus on equity and sustainability.

Notable progress has already been made. The stationary energy sector achieved an impressive 44% reduction in emissions since 1990, driven by the decarbonization of the electricity grid and advances in energy efficiency. The transportation sector saw a 3% reduction compared to 1990 levels, based on trends in data that we anticipate revising. While emissions from the waste sector are above the baseline, targeted programs to enhance recycling, divert waste from landfills, and engage the public are poised to address these challenges.

This report further highlights the City's success in decoupling emissions from economic growth, with emissions reduced and GDP still increasing. This progress illustrates that environmental sustainability and economic prosperity are not only compatible but mutually reinforcing as Los Angeles advances toward a greener, more resilient economy.

Looking ahead, the City will continue to innovate and implement bold climate strategies, guided by the Green New Deal's vision for carbon neutrality by 2050. LASAN, in partnership with the Mayor's Office of Energy and Sustainability, Climate Impact Team of the Office of the City Administrative Officer, and our City Departments, remains steadfast in refining the GHG inventory and developing policies that enable these ambitious goals. Los Angeles' journey toward sustainability is a testament to what can be accomplished through visionary leadership and collective action, serving as a beacon of environmental stewardship for cities worldwide.

5. Preparers

LA Sanitation & Environment (LASAN), recognized as a national leader in environmental services and programs, is a critical partner in the City's climate response and in advancing the path towards the City's climate goals. LASAN is committed to proactively addressing climate change and supporting climate action in line with our mission to protect public health and the environment.

Building on nearly a decade of experience, LASAN's Climate Action Program supports the City's path towards carbon neutrality as outlined by the Sustainable City pLAn and LA's Green New Deal. LASAN collaborates with City departments, policymakers, and outside agencies on climate-related reports and activities.

For more information about the Climate Action Program, please contact us at san.climateaction@lacity.org or visit us at www.lacitysan.org/climateaction.

Last updated: February 2026

Management

Doug Walters

Amanda Amaral

Climate Action Program

Daniel Brehm

Lily Carlson

Laura McAlister